arXiv:0902.4511v1 [cs.IT] 26 Feb 2009

Cyclic Codes and Sequences from Kasami-Welch Functions

February 26, 2009

Jinquan Luo San Ling and Chaoping Xing

---

[0] J.Luo is with the School of Mathematics, Yangzhou University, Jiangsu Province, 225009, China and with the Division of Mathematics, School of Physics and Mathematical Sciences, Nanyang Technological University, Singapore.

S.Ling and C.Xing are with the Division of Mathematics, School of Physics and Mathematical Sciences, Nanyang Technological University, Singapore.

E-mail addresses: jqluo@ntu.edu.sg, lingsan@ntu.edu.sg, xingcp@ntu.edu.sg.

Abstract

Let $q = 2^n$, $0 \leq k \leq n-1$ and $k \neq n/2$. In this paper we determine the value distribution of following exponential sums

$$\sum_{x \in \mathbb{F}_q} (-1)^{\mathrm{Tr}_1^n(\alpha x^{2^{3k}+1} + \beta x^{2^k+1})} \quad (\alpha, \beta \in \mathbb{F}_q)$$

and

$$\sum_{x \in \mathbb{F}_q} (-1)^{\mathrm{Tr}_1^n(\alpha x^{2^{3k}+1} + \beta x^{2^k+1} + \gamma x)} \quad (\alpha, \beta, \gamma \in \mathbb{F}_q)$$

where $\mathrm{Tr}_1^n : \mathbb{F}_{2^n} \to \mathbb{F}_2$ is the canonical trace mapping. As applications:

(1). We determine the weight distribution of the binary cyclic codes $\mathcal{C}_1$ and $\mathcal{C}_2$ with parity-check polynomials $h_2(x)h_3(x)$ and $h_1(x)h_2(x)h_3(x)$ respectively where $h_1(x)$, $h_2(x)$ and $h_3(x)$ are the minimal polynomials of π^{-1}, $\pi^{-(2^k+1)}$ and $\pi^{-(2^{3k}+1)}$ respectively for a primitive element π of $\mathbb{F}_q$.

(2). We determine the correlation distribution among a family of binary m-sequences.

Index terms: Exponential sum, Cyclic code, Sequence, Weight distribution, Correlation distribution

1 Introduction

Basic results on finite fields could be found in [19]. These notations are fixed throughout this paper except for specific statements.

- Let n be a positive integer, $q = 2^n$, $\mathbb{F}_q$ be the finite field of order q. Let π be a primitive element of $\mathbb{F}_q$.
- Let $\mathrm{Tr}_i^j : \mathbb{F}_{2^i} \rightarrow \mathbb{F}_{2^j}$ be the trace mapping, and $\chi(x) = (-1)^{\mathrm{Tr}_1^n(x)}$ be the canonical additive character on $\mathbb{F}_q$.
- Let k be a positive integer, $1 \leq k \leq n-1$ and $k \notin \{\frac{n}{4}, \frac{n}{2}, \frac{3n}{4}\}$. Let $d = \gcd(n, k)$, $q_0 = 2^d$, and $s = n/d$.
- Let $m = n/2$ (if n is even) and $\mu = (-1)^{m/d}$.

For binary cyclic code $\mathcal{C}$ with length l, let A_i be the number of codewords in $\mathcal{C}$ with Hamming weight i. The weight distribution $\{A_0, A_1, \cdots, A_l\}$ is an important research object for both theoretical and application interests in coding theory. Classical coding theory reveals that the weight of each codeword can be expressed by binary exponential sums so that the weight distribution of $\mathcal{C}$ can be determined if the corresponding exponential sums can be calculated explicitly (Kasami [16], [17], [18],van der Vlugt [28], Wolfmann[29]).

In general, let $q = 2^n$, $\mathcal{C}$ be the binary cyclic code with length $l = q - 1$ and parity-check polynomial

$$h(x) = h_1(x) \cdots h_u(x) \quad (u \geq 1)$$

where $h_i(x)$ $(1 \leq i \leq u)$ are distinct irreducible polynomials in $\mathbb{F}_2[x]$ with the same degree e_i $(1 \leq i \leq u)$, then $\dim_{\mathbb{F}_2}\mathcal{C} = \sum_{i=1}^{u} e_i$ .Let π^{-s_i} be a zero of $h_i(x)$, $1 \leq s_i \leq q-2$ $(1 \leq i \leq u)$. Then the codewords in $\mathcal{C}$ can be expressed by

$$c(\alpha_1, \cdots, \alpha_u) = (c_0, c_1, \cdots, c_{l-1}) \quad (\alpha_1, \cdots, \alpha_u \in \mathbb{F}_q)$$

where $c_i = \sum\limits_{\lambda=1}^{u} \mathrm{Tr}_1^n(\alpha_\lambda \pi^{is_\lambda})$ $(0 \le i \le l-1)$. Therefore the Hamming weight of the codeword $c = c(\alpha_1, \cdots, \alpha_u)$ is

$$\begin{aligned} w_H\,(c) &= \#\,\{i\,|0 \le i \le l-1, c_i \ne 0\} \\ &= l - \#\,\{i\,|0 \le i \le l-1, c_i = 0\} \\ &= l - \frac{1}{2}\sum_{i=0}^{l-1}\left(1 + (-1)^{\mathrm{Tr}_1^n\left(\sum\limits_{\lambda=1}^{u}\alpha_\lambda \pi^{is_\lambda}\right)}\right) \\ &= l - \frac{l}{2} - \frac{1}{2}\sum_{x\in\mathbb{F}_q^*}(-1)^{\mathrm{Tr}_1^n(f(x))} \\ &= \frac{l}{2} + \frac{1}{2} - \frac{1}{2}\,S(\alpha_1, \cdots, \alpha_u) \\ &= 2^{n-1} - \frac{1}{2}\,S(\alpha_1, \cdots, \alpha_u) \qquad (1) \end{aligned}$$

where $f(x) = \alpha_1 x^{s_1} + \alpha_2 x^{s_2} + \cdots + \alpha_u x^{s_u} \in \mathbb{F}_q[x]$, $\mathbb{F}_q^* = \mathbb{F}_q \backslash \{0\}$, and

$$S(\alpha_1, \cdots, \alpha_u) = \sum_{x\in\mathbb{F}_q}(-1)^{\mathrm{Tr}_1^n(\alpha_1 x^{s_1}+\cdots+\alpha_u x^{s_u})}.$$

In this way, the weight distribution of cyclic code $\mathcal{C}$ can be derived from the explicit evaluating of the exponential sums

$$S(\alpha_1, \cdots, \alpha_u) \quad (\alpha_1, \cdots, \alpha_u \in \mathbb{F}_q).$$

Let $h_1(x)$, $h_2(x)$ and $h_3(x)$ be the minimal polynomials of $\pi^{-1}, \pi^{-(2^k+1)}$ and $\pi^{-(2^{3k}+1)}$ over $\mathbb{F}_2$ respectively. Then $\deg h_i(x) = n$ for $i = 1, 2, 3$.

Let $\mathcal{C}_1$ and $\mathcal{C}_2$ be the binary cyclic codes with length $l = q-1$ and parity-check polynomials $h_2(x)h_3(x)$ and $h_1(x)h_2(x)h_3(x)$ respectively. It is a consequence that $\mathcal{C}_1(\mathcal{C}_2$, resp.) is the dual of the binary BCH code with designed distance 5 (7, resp.) whose zeroes include $\pi^{2^{3k}+1}$ and π^{2^k+1} ($\pi^{2^{3k}+1}$,π^{2^k+1} and π, resp.). Then we know that the dimensions of $\mathcal{C}_1$ and $\mathcal{C}_2$ are $2n$ and $3n$ respectively.

For $\alpha, \beta, \gamma \in \mathbb{F}_q$, define the exponential sums

$$T(\alpha, \beta) = \sum_{x\in\mathbb{F}_q}(-1)^{\mathrm{Tr}_1^n(\alpha x^{2^{3k}+1}+\beta x^{2^k+1})} \qquad (2)$$

and

$$S(\alpha,\beta,\gamma)=\sum_{x\in\mathbb{F}_q}(-1)^{\mathrm{Tr}_1^n(\alpha x^{2^{3k}+1}+\beta x^{2^k+1}+\gamma x)}. \tag{3}$$

Then the complete weight distributions of $\mathcal{C}_1$ and $\mathcal{C}_2$ can be derived from the explicit evaluation of $T(\alpha,\beta)$ and $S(\alpha,\beta,\gamma)$.

Another application of binary exponential sums is to obtain the cross correlation and auto-correlation distribution among binary sequences. Let $\mathcal{F}$ be a collection of binary m-sequences of period $q-1$ defined by

$$\mathcal{F}=\left\{\{a_i(t)\}_{i=0}^{q-2}\,|\,0\le i\le L-1\right\}$$

The *correlation function* of a_i and a_j for a shift τ is defined by

$$M_{i,j}(\tau)=\sum_{\lambda=0}^{q-2}(-1)^{a_i(\lambda)-a_j(\lambda+\tau)}\qquad\qquad(0\le\tau\le q-2).$$

Binary sequences with low cross correlation and auto-correlation are widely used in Code Division Multiple Access(CDMA) spread spectrum(see Paterson[23], Simon, Omura and Scholtz[26]). Pairs of binary m-sequences with few-valued auto and cross correlations have been extensively studied for several decades, see Canteaut, Charpin and Dobbertin [1], Cusick and Dobbertin [2], Ding, Helleseth and Lam[4], Ding, Helleseth and Martinsen[5], Dobbertin, Felke, Helleseth and Rosendahl [6], Gold [7], Helleseth [8],[9], Helleseth, Kholosha and Ness[10], Helleseth and Kumar [11], Hollmann and Xiang [12], Ness and Helleseth[20]-[21], Niho[22], Rosendahl[24], Yu and Gong[**?**]-[32] and references therein.

Define the collection of sequences

$$\mathcal{F}_1=\left\{a_{\alpha,\beta}=\left(a_{\alpha,\beta}(\pi^\lambda)\right)_{\lambda=0}^{q-2}\,|\,\alpha,\beta\in\mathbb{F}_q\right\}$$

where $a_{\alpha,\beta}(\pi^\lambda)=\mathrm{Tr}_1^n(\alpha\pi^{\lambda(2^{3k}+1)}+\beta\pi^{\lambda(2^k+1)}+\pi^\lambda)$.

If n/d is odd, define

$$\mathcal{F}_2=\left\{a_\beta=\left(a_\alpha(\pi^\lambda)\right)_{\lambda=0}^{q-2}\,|\,\beta\in\mathbb{F}_q\right\}\bigcup\left\{a=\left(a(\pi^\lambda)\right)_{\lambda=0}^{q-2}\right\}$$

where $a_\alpha(\pi^\lambda)=\mathrm{Tr}_1^n(\alpha\pi^{\lambda(2^{3k}+1)}+\pi^{\lambda(2^k+1)})$ and $a(\pi^\lambda)=\mathrm{Tr}_1^n(\pi^{\lambda(2^{3k}+1)})$.

In this correspondence we will study the following collection of m-sequences with period $q-1$

$$\mathcal{F}=\begin{cases}\mathcal{F}_1, & \text{if } n/d \text{ is even}\\ \mathcal{F}_1\cup\mathcal{F}_2, & \text{if } n/d \text{ is odd.}\end{cases} \tag{4}$$

This paper is presented as follows. In Section 2 we introduce some preliminaries and give auxiliary results. In Section 3 we will give the value distribution of $T(\alpha,\beta)$ for $\alpha\in\mathbb{F}_{2^m},\beta\in\mathbb{F}_q$ and the weight distribution of $\mathcal{C}_1$. In Section 3 we will determine the value distribution of $S(\alpha,\beta,\gamma)$, the correlation distribution among the sequences in $\mathcal{F}$, and then the weight distribution of $\mathcal{C}_2$. Most proofs of lemmas and theorems are presented in several appendices. The main tools are quadratic form theory over finite fields of characteristic 2, some moment identities on $T(\alpha,\beta)$ and a class of Artin-Schreier curves. This paper is binary analogue of [14] and [33].

2 Preliminaries

We follow the notations in Section 1. The first machinery to determine the values of exponential sums $T(\alpha,\beta)$ ($\alpha\in\mathbb{F}_{p^m},\beta\in\mathbb{F}_q$) defined in (2) is quadratic form theory over $\mathbb{F}_{q_0}$.

Let H be an $s\times s$ matrix over $\mathbb{F}_{q_0}$. For the quadratic form

$$F:\mathbb{F}_{q_0}^s\to\mathbb{F}_{q_0},\quad F(x)=XHX^T\quad (X=(x_1,\cdots,x_s)\in\mathbb{F}_{q_0}^s),\tag{5}$$

define r_F of F to be the rank of the skew-symmetric matrix $H+H^T$. Then r_F is even.

We have the following result on the exponential sum of binary quadratic forms.

***Lemma* 1.** *For the quadratic form $F=XHX^T$ defined in (5),*

$$\sum_{X\in\mathbb{F}_{q_0}^s}\zeta_p^{\mathrm{Tr}_1^d(F(X))}=\pm q_0^{s-\frac{r_F}{2}}\ \text{or}\ 0$$

Moreover, if $r_F=s$, then

$$\sum_{X\in\mathbb{F}_{q_0}^s}\zeta_p^{\mathrm{Tr}_1^d(F(X))}=\pm q_0^{\frac{s}{2}}$$

Proof. We can calculate

$$\begin{aligned}\left(\sum_{X\in\mathbb{F}_{q_0}^s}(-1)^{\mathrm{Tr}_1^d(F(X))}\right)^2&=\sum_{X,Z\in\mathbb{F}_{q_0}^s}(-1)^{\mathrm{Tr}_1^d\left(XHX^T+(X+Z)H(X+Z)^T\right)}\\&=\sum_{Z\in\mathbb{F}_{q_0}^s}(-1)^{\mathrm{Tr}_1^d(ZHZ^T)}\sum_{X\in\mathbb{F}_{q_0}^s}(-1)^{\mathrm{Tr}_1^d\left(Z(H+H^T)X^T\right)}\end{aligned}$$

The inner sum is zero unless $Z(H+H^T)=0$. Define

$$\mathcal{Z}=\left\{Z\in\mathbb{F}_{q_0}^s \mid Z(H+H^T)=0\right\}.$$

Then the map

$$\begin{aligned} \eta:\mathcal{Z} &\longrightarrow \mathbb{F}_2 \\ Z &\mapsto \mathrm{Tr}_1^d(ZHZ^T) \end{aligned} \tag{6}$$

is an additive group homomorphism.

If η is surjective, then there are exactly one half $z\in\mathcal{Z}$ mapping to 0 and 1 respectively. Hence $\sum\limits_{X\in\mathbb{F}_{q_0}^s}(-1)^{\mathrm{Tr}_1^d(F(X))}=0$. Otherwise $\mathrm{Im}(\eta)=\{0\}$ and $\left(\sum\limits_{X\in\mathbb{F}_{q_0}^s}(-1)^{\mathrm{Tr}_1^d(F(X))}\right)^2=q_0^s\cdot q_0^{s-r_F}$. Hence $\sum\limits_{X\in\mathbb{F}_{q_0}^s}(-1)^{\mathrm{Tr}_1^d(F(X))}=\pm q_0^{s-\frac{r_F}{2}}$.

If $r_F=s$, then $\mathcal{Z}=\{0\}$ and $\mathrm{Im}(\eta)=\{0\}$. Therefore $\sum\limits_{X\in\mathbb{F}_{q_0}^s}(-1)^{\mathrm{Tr}_1^d(F(X))}=\pm q_0^{\frac{s}{2}}$.

□

The following result, which has been proven in [19], Chap. 6, will be used in Section 4.

Lemma 2. *For the fixed quadratic form defined in (5), the value distribution of* $\sum\limits_{X\in\mathbb{F}_{q_0}^s}(-1)^{\mathrm{Tr}_1^d(F(X)+AX^T)}$ *when* A *runs through* $\mathbb{F}_{q_0}^s$ *is shown as following*

value	*multiplicity*
0	$q_0^s-q_0^{r_F}$
$q_0^{s-\frac{r_F}{2}}$	$\frac{1}{2}(q_0^{r_F}+q_0^{\frac{r_F}{2}})$
$-q_0^{s-\frac{r_F}{2}}$	$\frac{1}{2}(q_0^{r_F}-q_0^{\frac{r_F}{2}})$

Since $s=n/d$, the field $\mathbb{F}_q$ is a vector space over $\mathbb{F}_{q_0}$ with dimension s. We fix a basis $v_1,\cdots,v_s$ of $\mathbb{F}_q$ over $\mathbb{F}_{q_0}$. Then each $x\in\mathbb{F}_q$ can be uniquely expressed as

$$x=x_1v_1+\cdots+x_sv_s \quad (x_i\in\mathbb{F}_{q_0}).$$

Thus we have the following $\mathbb{F}_{q_0}$-linear isomorphism:

$$\mathbb{F}_q\xrightarrow{\sim}\mathbb{F}_{q_0}^s,\quad x=x_1v_1+\cdots+x_sv_s\mapsto X=(x_1,\cdots,x_s).$$

With this isomorphism, a function $f : \mathbb{F}_q \to \mathbb{F}_{q_0}$ induces a function $F : \mathbb{F}_{q_0}^s \to \mathbb{F}_{q_0}$ where for $X = (x_1, \cdots, x_s) \in \mathbb{F}_{q_0}^s, F(X) = f(x)$ with $x = x_1 v_1 + \cdots + x_s v_s$. In this way, function $f(x) = \mathrm{Tr}_d^n(\gamma x)$ for $\gamma \in \mathbb{F}_q$ induces a linear form

$$F(X) = \mathrm{Tr}_d^n(\gamma x) = \sum_{i=1}^{s} \mathrm{Tr}_d^n(\gamma v_i) x_i = A_\gamma X^T \tag{7}$$

where $A_\gamma = \left(\mathrm{Tr}_d^n(\gamma v_1), \cdots, \mathrm{Tr}_d^n(\gamma v_s)\right)$, and $f_{\alpha,\beta}(x) = \mathrm{Tr}_d^n(\alpha x^{p^{3k}+1} + \beta x^{p^k+1})$ for $\alpha, \beta \in \mathbb{F}_q$ induces a quadratic form

$$F_{\alpha,\beta}(X) = X H_{\alpha,\beta} X^T \tag{8}$$

From Lemma 1, for $\alpha, \beta, \gamma \in \mathbb{F}_q$, in order to determine the values of

$$T(\alpha, \beta) = \sum_{x \in \mathbb{F}_q} (-1)^{\mathrm{Tr}_1^n(\alpha x^{2^{3k}+1} + \beta x^{2^k+1})} = \sum_{X \in \mathbb{F}_{q_0}^s} (-1)^{\mathrm{Tr}_1^d\left(X H_{\alpha,\beta} X^T\right)}$$

and

$$S(\alpha, \beta, \gamma) = \sum_{x \in \mathbb{F}_q} (-1)^{\mathrm{Tr}_1^n(\alpha x^{2^{3k}+1} + \beta x^{2^k+1} + \gamma x)} = \sum_{X \in \mathbb{F}_{q_0}^s} (-1)^{\mathrm{Tr}_1^d\left(X H_{\alpha,\beta} X^T + A_\gamma X^T\right)},$$

we need to determine the rank of $H_{\alpha,\beta} + H_{\alpha,\beta}^T$ over $\mathbb{F}_{q_0}$.

Define $d' = \gcd(n, 2k)$. Then an easy observation shows

$$d' = \begin{cases} d, & \text{if } n/d \text{ is odd;} \\ 2d, & \text{otherwise.} \end{cases} \tag{9}$$

***Lemma* 3.** *For $(\alpha, \beta) \in \mathbb{F}_{p^m} \times \mathbb{F}_q \backslash \{(0,0)\}$, let $r_{\alpha,\beta}$ be the rank of $H_{\alpha,\beta} + H_{\alpha,\beta}^T$. Then we have*

(i). if n/d is odd, then then the possible values of $r_{\alpha,\beta}$ are $s-1$ and $s-3$.

(ii). if n/d is even, then the possible values of $r_{\alpha,\beta}$ are s, $s-2$, $s-4$ and $s-6$.

In order to determine the value distribution of $T(\alpha, \beta)$ for $\alpha, \beta \in \mathbb{F}_q$, we need the following result on moments of $T(\alpha, \beta)$ and $S(\alpha, \beta, \gamma)$.

***Lemma* 4.** *For the exponential sum $T(\alpha,\beta)$ and $S(\alpha,\beta,\gamma)$,*

$$(i).\quad \sum_{\alpha,\beta\in\mathbb{F}_q} T(\alpha,\beta) = 2^{2n};$$

(ii). if n/d is even, then

$$\sum_{\alpha,\beta\in\mathbb{F}_q} T(\alpha,\beta)^2 = (2^{n+d} + 2^n - 2^d)\cdot 2^{2n}$$

(iii). if n/d is even, then

$$\sum_{\alpha,\beta\in\mathbb{F}_q} T(\alpha,\beta)^3 = (2^{n+3d} + 2^n - 2^{3d})\cdot 2^{2n}.$$

$$(iv).\quad \sum_{\alpha,\beta,\gamma\in\mathbb{F}_q} S(\alpha,\beta,\gamma)^3 = (2^{n+d} + 2^n - 2^d)\cdot 2^{3n}.$$

Proof. see **Appendix A.** □

***Remark*.** We could calculate the moment identities $\sum_{\alpha,\beta\in\mathbb{F}_q} T(\alpha,\beta)^i$ and $\sum_{\alpha,\beta,\gamma\in\mathbb{F}_q} T(\alpha,\beta,\gamma)^i$ for $1 \le i \le 3$. But the result shown above is enough to determine the value distributions of $T(\alpha,\beta)$ and $S(\alpha,\beta,\gamma)$.

In the case n/d is even, we could determine the explicit values of $T(\alpha,\beta)$. To this end we will study a class of Artin-Schreier curves. A similar technique has been applied in Coulter [3], Theorem 6.1.

***Lemma* 5.** *Suppose $(\alpha,\beta) \in (\mathbb{F}_q \times \mathbb{F}_q)\backslash\{0,0\}$ and $d' = 2d$. Let N be the number of $\mathbb{F}_q$-rational (affine) points on the curve*

$$\alpha x^{2^{3k}+1} + \beta x^{2^k+1} = y^{2^d} + y. \tag{10}$$

Then

$$N = q + (2^d - 1)\cdot T(\alpha,\beta).$$

Proof. see **Appendix A.** □

Now we give an explicit evaluation of $T(\alpha,\beta)$ in the case n/d is even.

***Lemma* 6.** *Assumptions as in Lemma 5 and let $n = 2m$, then*

$$T(\alpha,\beta) = \begin{cases} \mu 2^m, & \text{if } r_{\alpha,\beta} = s \\ -\mu 2^{m+d}, & \text{if } r_{\alpha,\beta} = s-2 \\ \mu 2^{m+2d}, & \text{if } r_{\alpha,\beta} = s-4 \\ -\mu 2^{m+3d}, & \text{if } r_{\alpha,\beta} = s-6. \end{cases}$$

where $\mu = (-1)^{m/d}$.

Proof. Consider the $\mathbb{F}_q$-rational (affine) points on the Artin-Schreier curve in Lemma 5. It is easy to verify that $(0, y)$ with $y \in \mathbb{F}_{2^d}$ are exactly the points on the curve with $x = 0$. If (x, y) with $x \neq 0$ is a point on this curve, then so are $(tx, t^{2^d+1}y)$ with $t^{2^{2d}-1} = 1$ (note that $2^{3k} + 1 \equiv 2^k + 1 \equiv 2^d + 1 \pmod{2^{2d} - 1}$ since $3k/d$ and k/d are both odd by (9)). In total, we have

$$q + (2^d - 1)T(\alpha, \beta) = N \equiv 2^d \pmod{2^{2d} - 1}$$

which yields

$$T(\alpha, \beta) \equiv 1 \pmod{2^d + 1}.$$

We only consider the case $r_{\alpha,\beta} = s$ and m/d is odd. The other cases are similar. In this case $T(\alpha, \beta) = \pm 2^m$. Assume $T(\alpha, \beta) = 2^m$. Then $2^d + 1 \mid 2^m - 1$ which contradicts to m/d is odd. Therefore $T(\alpha, \beta) = -2^m$. $\square$

Remark. (i) In the case n/d is even, the non-binary case (see [14], Lemma 6) and the binary case could be solved in a unified way. But the case n/d odd is quite different.

(ii). Applying Lemma 6 to Lemma 5, we could determine the number of rational points on the curve (10).

3 Exponential Sums $T(\alpha, \beta)$ and Cyclic Code $\mathcal{C}_1$

Define

$$N_i = \{(\alpha, \beta) \in \mathbb{F}_q \times \mathbb{F}_q \backslash \{(0, 0)\} \,| r_{\alpha,\beta} = s - i\}.$$

Then $n_i = |N_i|$.

According to the possible values of $T(\alpha, \beta)$ given by Lemma 1 (setting $F(X) = XH_{\alpha,\beta}X^T = \mathrm{Tr}_d^n(\alpha x^{2^{3k}+1} + \beta x^{2^k+1})$), we define that for $\varepsilon = \pm 1$,

$$N_{\varepsilon,i} = \left\{(\alpha, \beta) \in \mathbb{F}_q^2 \backslash \{(0, 0)\} \,\middle|\, T(\alpha, \beta) = \varepsilon 2^{\frac{n+id}{2}}\right\}$$

and $n_{\varepsilon,i} = |N_{\varepsilon,i}|$.

Now we could give the value distribution of $T(\alpha, \beta)$ and the weight distribution of $\mathcal{C}_1$. We will only focus on the case n/d is even since the odd case has been solved in [12].

***Theorem* 1.** *The value distribution of the multi-set $\{T(\alpha, \beta) \,| \alpha, \beta \in \mathbb{F}_q\}$ and the weight distribution of $\mathcal{C}_1$ are shown as following (Column 1 is the value of $T(\alpha, \beta)$,* *Column 2 is the weight of* $c(\alpha,\beta) = \mathrm{Tr}_1^n\left(\alpha\pi^{i(2^{3k}+1)} + \beta\pi^{i(2^k+1)}\right)_{i=0}^{q-2}$ *and Column 3 is the corresponding multiplicity).*

(i). For the case n/d *is odd,*

values	*weights*	*multiplicity*
$2^{(n+d)/2}$	$2^{n-1} - 2^{(n+d-2)/2}$	$\left(2^{n-d-1} + 2^{(n-d-2)/2}\right)(2^n - 1)$
$-2^{(n+d)/2}$	$2^{n-1} + 2^{(n+d-2)/2}$	$\left(2^{n-d-1} - 2^{(n-d-2)/2}\right)(2^n - 1)$
0	2^{n-1}	$\left(2^n - 2^{n-d} + 1\right)(2^n - 1)$
2^n	0	1

(ii). For the case n/d *is even,*

values	*weights*	*multiplicity*
$\mu 2^m$	$2^{n-1} - \mu 2^{m-1}$	$\frac{(2^n-1)\left(2^{n+6d}-2^{n+4d}-2^{n+d}+\mu 2^{m+5d}-\mu 2^{m+4d}+2^{6d}\right)}{(2^d+1)(2^{2d}-1)(2^{3d}+1)}$
$-\mu 2^{m+d}$	$2^{n-1} + \mu 2^{m+d-1}$	$\frac{(2^n-1)(2^{n+3d}+2^{n+2d}-2^n-2^{n-d}-2^{n-2d}-\mu 2^{m+3d}+\mu 2^m+2^{3d})}{(2^d+1)^2(2^{2d}-1)}$
$\mu 2^{m+2d}$	$2^{n-1} - \mu 2^{m+2d-1}$	$\frac{(2^{m-d}+\mu)(2^{m+d}+2^m-2^{m-2d}-\mu 2^d)(2^n-1)}{(2^d+1)^3(2^d-1)}$
$-\mu 2^{m+3d}$	$2^{n-1} + \mu 2^{m+3d-1}$	$\frac{(2^{m-2d}-\mu)(2^{m-d}+\mu)(2^n-1)}{(2^d+1)(2^{2d}-1)(2^{3d}+1)}$
2^n	0	1

where $\mu = (-1)^{m/d}$.

Proof. see [12] for (i) and **Appendix B** for (ii). ☐

Remark. (i). In the case $k \in \left\{\frac{n}{4}, \frac{n}{2}, \frac{3n}{4}\right\}$, the exponential sum $T(\alpha,\beta) = \sum_{x\in\mathbb{F}_q}(-1)^{\mathrm{Tr}_1^n\left(\left(\alpha^{2^k}+\beta\right)x^{2^k+1}\right)}$ and the cyclic code $\mathcal{C}_1$ has been extensively studied, for example, see [3], [7].

(ii). In the case $k \in \left\{\frac{n}{6}, \frac{5n}{6}\right\}$, the exponential sum $T(\alpha,\beta)$ and the weight distribution of $\mathcal{C}_1$ has shown in [15]. In the case $k \in \{\frac{n}{3}, \frac{2n}{3}\}$, the exponential sum $T(\alpha,\beta) = \sum_{x\in\mathbb{F}_q}(-1)^{\mathrm{Tr}_1^n\left(\alpha^{2^{n-1}}x+\beta x^{2^k+1}\right)}$ is a special case in [3], [7].

(iii). In the case n/d is even. Since $\gcd(2^n-1, 2^k+1) = 2^d+1$, the first $l' = \frac{2^n-1}{2^d+1}$ coordinates of each codeword of $\mathcal{C}_1$ form a cyclic code $\mathcal{C}_1'$ with length l' and dimension $2n$. Let $(A_0', \cdots, A_{l'}')$ be the weight distribution of $\mathcal{C}_1'$, then $A_i' = A_{(2^d+1)i}$ $(0 \le i \le l')$.

4 Results on Correlation Distribution of Sequences and Cyclic Code $\mathcal{C}_2$

Recall $\phi_{\alpha,\beta}(x)$ in the proof of Lemma 3 and $N_{i,\varepsilon}$ in the proof of Theorem 1. Finally we will determine the value distribution of $S(\alpha,\beta,\gamma)$, the correlation distribution among sequences in $\mathcal{F}$ defined in (4) and the weight distribution of $\mathcal{C}_2$ defined in Section 1.

Theorem 2. *The value distribution of the multi-set* $\{S(\alpha,\beta,\gamma)\,|\alpha,\beta,\gamma \in \mathbb{F}_q\}$ *and the weight distribution of* $\mathcal{C}_2$ *are shown as following (Column 1 is the value of* $S(\alpha,\beta,\gamma)$*, Column 2 is the weight of* $c(\alpha,\beta,\gamma) = \left(\mathrm{Tr}_1^n(\alpha\pi^{i(2^{3k}+1)} + \beta\pi^{i(2^k+1)} + \gamma\pi^i)\right)_{i=0}^{q-2}$ *and Column 3 is the corresponding multiplicity).*

(i). For the case n/d *is odd (that is,* $d' = d$*),*

values	*weights*	*multiplicity*
$2^{(n+d)/2}$	$2^{n-1} - 2^{(n+d-2)/2}$	$\frac{(2^{n-d-1}+2^{(n-d-2)/2})(2^n-1)(2^{n+2d}-2^n-2^{n-d}+2^{2d})}{2^{2d}-1}$
$-2^{(n+d)/2}$	$2^{n-1} + 2^{(n+d-2)/2}$	$\frac{(2^{n-d-1}-2^{(n-d-2)/2})(2^n-1)(2^{n+2d}-2^n-2^{n-d}+2^{2d})}{2^{2d}-1}$
$2^{(n+3d)/2}$	$2^{n-1} - 2^{(n+3d-2)/2}$	$\frac{(2^{n-3d-1}+2^{(n-3d-2)/2})(2^{n-d}-1)(2^n-1)}{2^{2d}-1}$
$-2^{(n+3d)/2}$	$2^{n-1} + 2^{(n+3d-2)/2}$	$\frac{(2^{n-3d-1}-2^{(n-3d-2)/2})(2^{n-d}-1)(2^n-1)}{2^{2d}-1}$
0	2^{n-1}	$(2^n-1)(2^{2n}-2^{2n-d}+2^{2n-4d}+2^n-2^{n-d}-2^{n-3d}+1)$
2^n	0	1

(ii). For the case n/d *is even,*

values	*weights*	*multiplicity*
2^m	$2^{n-1}-2^{m-1}$	$\frac{(2^{n-1}+2^{m-1})(2^n-1)\left(2^{n+6d}-2^{n+4d}-2^{n+d}+\mu 2^{m+5d}-\mu 2^{m+4d}+2^{6d}\right)}{(2^d+1)(2^{2d}-1)(2^{3d}+1)}$
-2^m	$2^{n-1}+2^{m-1}$	$\frac{(2^{n-1}-2^{m-1})(2^n-1)\left(2^{n+6d}-2^{n+4d}-2^{n+d}+\mu 2^{m+5d}-\mu 2^{m+4d}+2^{6d}\right)}{(2^d+1)(2^{2d}-1)(2^{3d}+1)}$
2^{m+d}	$2^{n-1}-2^{m+d-1}$	$\frac{(2^{n-2d-1}+2^{m-d-1})(2^n-1)(2^{n+3d}+2^{n+2d}-2^n-2^{n-d}-2^{n-2d}-\mu 2^{m+3d}+\mu 2^m+2^{3d})}{(2^d+1)^2(2^{2d}-1)}$
-2^{m+d}	$2^{n-1}+2^{m+d-1}$	$\frac{(2^{n-2d-1}-2^{m-d-1})(2^n-1)(2^{n+3d}+2^{n+2d}-2^n-2^{n-d}-2^{n-2d}-\mu 2^{m+3d}+\mu 2^m+2^{3d})}{(2^d+1)^2(2^{2d}-1)}$
2^{m+2d}	$2^{n-1}-2^{m+2d-1}$	$\frac{(2^{m-d}+\mu)(2^{m+d}+2^m-2^{m-2d}-\mu 2^d)(2^{n-4d-1}+2^{m-2d-1})(2^n-1)}{(2^d+1)^3(2^d-1)}$
-2^{m+2d}	$2^{n-1}+2^{m+2d-1}$	$\frac{(2^{m-d}+\mu)(2^{m+d}+2^m-2^{m-2d}-\mu 2^d)(2^{n-4d-1}-2^{m-2d-1})(2^n-1)}{(2^d+1)^3(2^d-1)}$
2^{m+3d}	$2^{n-1}-2^{m+3d-1}$	$\frac{(2^{m-2d}-\mu)(2^{m-d}+\mu)(2^{n-6d-1}+2^{m-3d-1})(2^n-1)}{(2^d+1)(2^{2d}-1)(2^{3d}+1)}$
-2^{m+3d}	$2^{n-1}+2^{m+3d-1}$	$\frac{(2^{m-2d}-\mu)(2^{m-d}+\mu)(2^{n-6d-1}-2^{m-3d-1})(2^n-1)}{(2^d+1)(2^{2d}-1)(2^{3d}+1)}$
0	2^{n-1}	$\frac{(2^n-1)(2^{2n}+2^{2n-9d}-\varepsilon(2^{3m}-2^{3m-d}-2^{3m-3d}+2^{3m-5d}+2^{3m-7d}-2^{3m-8d})+2^n-2^{n-d}-2^{n-4d}-2^{n-6d}+2^d+1)}{2^d+1}$
2^n	0	1

where $\mu=(-1)^{m/d}$.

Proof. See **Appendix C.** ☐

In order to give the correlation distribution among the sequences in $\mathcal{F}$, we need an easy observation.

Lemma 7. *For any given* $\gamma\in\mathbb{F}_q^*$, *when* (α,β) *runs through* $\mathbb{F}_q\times\mathbb{F}_q$, *the distribution of* $S(\alpha,\beta,\gamma)$ *is the same as* $S(\alpha,\beta,1)$.

As a consequence of Theorem 1, Theorem 2 and Lemma 7, we could give the correlation distribution amidst the sequences in $\mathcal{F}$.

Theorem 3. *Let* $1\le k\le n-1$ *and* $k\notin\left\{\frac{n}{6},\frac{n}{4},\frac{n}{2},\frac{3n}{4},\frac{5n}{6}\right\}$. *The collection* $\mathcal{F}$ *defined in (4) is a family of* 2^{2n} *(*n/d *is odd) or* $2^{2n}+2^n+1$ *(*n/d *is even) binary sequences with period* $q-1$. *Its correlation distribution is given as follows.*

(i). For the case n/d *is odd (that is,* $d'=d$*),*

values	*multiplicity*
$2^{(n+d)/2}-1$	$\frac{(2^{n-d-1}+2^{(n-d-2)/2})(2^{n+2d}-2^{n}-2^{n-d}+2^{2d})(2^{3n}-2^{n+1})}{2^{2d}-1}$
$-2^{(n+d)/2}-1$	$\frac{(2^{n-d-1}-2^{(n-d-2)/2})(2^{n+2d}-2^{n}-2^{n-d}+2^{2d})(2^{3n}-2^{n+1})}{2^{2d}-1}$
$2^{(n+3d)/2}-1$	$\frac{(2^{n-3d-1}+2^{(n-3d-2)/2})(2^{n-d}-1)(2^{3n}-2^{n+1})}{2^{2d}-1}$
$-2^{(n+3d)/2}-1$	$\frac{(2^{n-3d-1}-2^{(n-3d-2)/2})(2^{n-d}-1)(2^{3n}-2^{n+1})}{2^{2d}-1}$
-1	$(2^{2n}-2^{2n-d}+2^{2n-4d}+2^{n}-2^{n-d}-2^{n-3d}+1)(2^{3n}-2^{n+1})+2^{n}$
$2^{n}-1$	$2^{2n}+2^{n}$

(ii). For the case $n/d \equiv 0 \pmod 4$.

values	*multiplicity*
$2^{m}-1$	$\frac{2^{2n}\left(2^{n+6d}-2^{n+4d}-2^{n+d}+2^{m+5d}-2^{m+4d}+2^{6d}\right)(2^{2n-1}+2^{3m-1}-2^{n}-2^{m}+1)}{(2^{d}+1)(2^{2d}-1)(2^{3d}+1)}$
$-2^{m}-1$	$\frac{2^{2n}(2^{n-1}-2^{m-1})(2^{n}-2)\left(2^{n+6d}-2^{n+4d}-2^{n+d}+2^{m+5d}-2^{m+4d}+2^{6d}\right)}{(2^{d}+1)(2^{2d}-1)(2^{3d}+1)}$
$2^{m+d}-1$	$\frac{2^{2n}(2^{n-2d-1}+2^{m-d-1})(2^{n}-2)(2^{n+3d}+2^{n+2d}-2^{n}-2^{n-d}-2^{n-2d}-2^{m+3d}+2^{m}+2^{3d})}{(2^{d}+1)^{2}(2^{2d}-1)}$
$-2^{m+d}-1$	$\frac{2^{2n}(2^{n+3d}+2^{n+2d}-2^{n}-2^{n-d}-2^{n-2d}-2^{m+3d}+2^{m}+2^{3d})(2^{2n-2d-1}-2^{3m-d-1}-2^{n-2d}+2^{m-d}+1)}{(2^{d}+1)^{2}(2^{2d}-1)}$
$2^{m+2d}-1$	$\frac{2^{2n}(2^{m-d}+1)(2^{m+d}+2^{m}-2^{m-2d}-2^{d})(2^{2n-4d-1}+2^{3m-2d-1}-2^{n-4d}-2^{m-2d}+1)}{(2^{d}+1)^{3}(2^{d}-1)}$
$-2^{m+2d}-1$	$\frac{2^{2n}(2^{m-d}+\mu)(2^{m+d}+2^{m}-2^{m-2d}-\mu 2^{d})(2^{n-4d-1}-2^{m-2d-1})(2^{n}-2)}{(2^{d}+1)^{3}(2^{d}-1)}$
$2^{m+3d}-1$	$\frac{2^{2n}(2^{m-2d}-1)(2^{m-d}+1)(2^{n-6d-1}+2^{m-3d-1})(2^{n}-2)}{(2^{d}+1)(2^{2d}-1)(2^{3d}+1)}$
$-2^{m+3d}-1$	$\frac{2^{2n}(2^{m-2d}-1)(2^{m-d}+1)(2^{2n-6d-1}-2^{3m-3d-1}-2^{n-6d}+2^{m-3d}+1)}{(2^{d}+1)(2^{2d}-1)(2^{3d}+1)}$
-1	$\frac{2^{2n}(2^{n}-2)\left(2^{2n}+2^{2n-9d}-2^{3m}+2^{3m-d}+2^{3m-3d}-2^{3m-5d}-2^{3m-7d}+2^{3m-8d}+2^{n}-2^{n-d}-2^{n-4d}-2^{n-6d}+2^{d}+1\right)}{2^{d}+1}$
$2^{n}-1$	2^{2n}

(iii). For the case $n/d \equiv 2 \pmod 4$,

values	*multiplicity*
2^m-1	$\frac{2^{2n}(2^{n-1}+2^{m-1})(2^n-2)\left(2^{n+6d}-2^{n+4d}-2^{n+d}-2^{m+5d}+2^{m+4d}+2^{6d}\right)}{(2^d+1)(2^{2d}-1)(2^{3d}+1)}$
-2^m-1	$\frac{2^{2n}\left(2^{n+6d}-2^{n+4d}-2^{n+d}-2^{m+5d}+2^{m+4d}+2^{6d}\right)(2^{2n-1}-2^{3m-1}-2^n+2^m+1)}{(2^d+1)(2^{2d}-1)(2^{3d}+1)}$
$2^{m+d}-1$	$\frac{2^{2n}(2^{n+3d}+2^{n+2d}-2^n-2^{n-d}-2^{n-2d}+2^{m+3d}-2^m+2^{3d})(2^{2n-2d-1}+2^{3m-d-1}-2^{n-2d}-2^{m-d}+1)}{(2^d+1)^2(2^{2d}-1)}$
$-2^{m+d}-1$	$\frac{2^{2n}(2^{n-2d-1}-2^{m-d-1})(2^n-2)(2^{n+3d}+2^{n+2d}-2^n-2^{n-d}-2^{n-2d}+2^{m+3d}-2^m+2^{3d})}{(2^d+1)^2(2^{2d}-1)}$
$2^{m+2d}-1$	$\frac{2^{2n}(2^{m-d}-1)(2^{m+d}+2^m-2^{m-2d}+2^d)(2^{n-4d-1}+2^{m-2d-1})(2^n-2)}{(2^d+1)^3(2^d-1)}$
$-2^{m+2d}-1$	$\frac{2^{2n}(2^{m-d}-1)(2^{m+d}+2^m-2^{m-2d}+2^d)(2^{2n-4d-1}-2^{3m-2d-1}-2^{n-4d}+2^{m-2d}+1)}{(2^d+1)^3(2^d-1)}$
$2^{m+3d}-1$	$\frac{2^{2n}(2^{m-2d}+1)(2^{m-d}-1)(2^{2n-6d-1}-2^{3m-3d-1}-2^{n-6d}+2^{m-3d}+1)}{(2^d+1)(2^{2d}-1)(2^{3d}+1)}$
$-2^{m+3d}-1$	$\frac{2^{2n}(2^{m-2d}+1)(2^{m-d}-1)(2^{n-6d-1}-2^{m-3d-1})(2^n-2)}{(2^d+1)(2^{2d}-1)(2^{3d}+1)}$
-1	$\frac{2^{2n}(2^n-2)\left(2^{2n}+2^{2n-9d}+2^{3m}-2^{3m-d}-2^{3m-3d}+2^{3m-5d}+2^{3m-7d}-2^{3m-8d}+2^n-2^{n-d}-2^{n-4d}-2^{n-6d}+2^d+1\right)}{2^d+1}$
2^n-1	2^{2n}

Proof. see **Appendix C.** □

Table 1: **Some Sequence Families with Low Correlation**

sequence	n	period	family size	$C_{\max}$
Gold[7]	odd	2^n-1	2^n+1	$2^{(n+1)/2}+1$
Rothaus[25]	even	2^n-1	$2^{2n}+2^n+1$	$2^{(n+3)/2}+1$
Yu and Gong[31], $1<\rho\leq\frac{n-1}{2}$	odd	2^n-1	$2^{n\rho}$	$2^{(n+2\rho-1)/2}+1$
Generalized Kasami(Large Set)[18],[30]	even	2^n-1	$2^{3n/2}+1$	$2^{(n+2)/2}$
$\mathcal{F}$ with $d=1$	odd	2^n-1	$2^{2n}+2^n+1$	$2^{(n+3)/2}+1$

5 Appendix A

Proof of Lemma 3:

For $Y = (y_1, \cdots, y_s) \in \mathbb{F}_{q_0}^s$, $y = y_1 v_1 + \cdots + y_s v_s \in \mathbb{F}_q$, we know that

$$F_{\alpha,\beta}(X+Y) - F_{\alpha,\beta}(X) - F_{\alpha,\beta}(Y) = 2XH_{\alpha,\beta}Y^T \tag{11}$$

is equal to

$$f_{\alpha,\beta}(x+y) - f_{\alpha,\beta}(x) - f_{\alpha,\beta}(y) = \mathrm{Tr}_d^n \left(y^{2^{3k}} (\alpha^{2^{3k}} x^{2^{6k}} + \beta^{2^{3k}} x^{2^{4k}} + \beta^{2^{2k}} x^{2^{2k}} + \alpha x) \right) \tag{12}$$

Let

$$\phi_{\alpha,\beta}(x) = \alpha^{2^{3k}} x^{2^{6k}} + \beta^{2^{3k}} x^{2^{4k}} + \beta^{2^{2k}} x^{2^{2k}} + \alpha x. \tag{13}$$

Therefore,

$$\begin{aligned} r_{\alpha,\beta} = r &\Leftrightarrow \text{the number of common solutions of } XH_{\alpha,\beta}Y^T = 0 \text{ for all } Y \in \mathbb{F}_{q_0}^s \text{ is } q_0^{s-r}, \\ &\Leftrightarrow \text{the number of common solutions of } \mathrm{Tr}_d^n \left(y^{2^{3k}} \cdot \phi_{\alpha,\beta}(x) \right) = 0 \text{ for all } y \in \mathbb{F}_q \text{ is } q_0^{s-r}, \\ &\Leftrightarrow \phi_{\alpha,\beta}(x) = 0 \text{ has } q_0^{s-r} \text{ solutions in } \mathbb{F}_q. \end{aligned}$$

For a fixed algebraic closure $\mathbb{F}_{2^\infty}$ of $\mathbb{F}_2$, since the degree of 2^{2k}-linearized polynomial $\phi_{\alpha,\beta}(x)$ is 2^{6k} and $\phi_{\alpha,\beta}(x) = 0$ has no multiple roots in $\mathbb{F}_{2^\infty}$ (this fact follows from $\phi'_{\alpha,\beta}(x) = \alpha \in \mathbb{F}_q^*$), then the zeroes of $\phi_{\alpha,\beta}(x)$ in $\mathbb{F}_{2^\infty}$, say V, form an $\mathbb{F}_{2^{2k}}$-vector space of dimension 3. Note that $\gcd(n, 2k) = d'$. Then $V \cap \mathbb{F}_{2^n}$ is a vector space on $\mathbb{F}_{2^{\gcd(n,2k)}} = \mathbb{F}_{2^{d'}}$ of dimension less that or equal to 3 since any elements in $\mathbb{F}_{2^n}$ which are linear independent over $\mathbb{F}_{2^{d'}}$ are also linear independent over $\mathbb{F}_{2^{2k}}$(see [27], Lemma 4).

(i). For the case n/d is odd, then $d' = d$ and $r_{\alpha,\beta} \geq s - 3$. Note that $r_{\alpha,\beta}$ must be even. Hence the possible values of $r_{\alpha,\beta}$ are $s-1$ and $s-3$.

(ii). For the case n/d is even, then $d' = 2d$ and the possible values of $r_{\alpha,\beta}$ are s, $s-2$, $s-4$ and $s-6$.

□

Proof of Lemma 4: (i). We observe that

$$\begin{aligned} \sum_{\alpha,\beta \in \mathbb{F}_q} T(\alpha, \beta) &= \sum_{\alpha,\beta \in \mathbb{F}_q} \sum_{x \in \mathbb{F}_q} (-1)^{\mathrm{Tr}_1^n(\alpha x^{2^{3k}+1} + \beta x^{2^k+1})} \\ &= \sum_{x \in \mathbb{F}_q} \sum_{\alpha \in \mathbb{F}_q} (-1)^{\mathrm{Tr}_1^n(\alpha x^{2^{3k}+1})} \sum_{\beta \in \mathbb{F}_q} (-1)^{\mathrm{Tr}_1^n(\beta x^{2^k+1})} = q \cdot \sum_{\substack{\alpha \in \mathbb{F}_q \\ x=0}} (-1)^{\mathrm{Tr}_1^n(\alpha x^{2^{3k}+1})} = 2^{2n}. \end{aligned}$$

(ii). We can calculate

$$\sum_{\alpha,\beta\in\mathbb{F}_q} T(\alpha,\beta)^2 = \sum_{x,y\in\mathbb{F}_q}\sum_{\alpha\in\mathbb{F}_q}(-1)^{\mathrm{Tr}_1^n\left(\alpha\left(x^{2^{3k}+1}+y^{2^{3k}+1}\right)\right)}\sum_{\beta\in\mathbb{F}_q}(-1)^{\mathrm{Tr}_1^n\left(\beta\left(x^{2^k+1}+y^{2^k+1}\right)\right)}$$
$$= M_2\cdot 2^{2n}$$

where M_2 is the number of solutions to the equation

$$\begin{cases} x^{2^{3k}+1}+y^{2^{3k}+1}=0 \\ x^{2^k+1}+y^{2^k+1}=0 \end{cases} \tag{14}$$

If $xy=0$ satisfying (14), then $x=y=0$. Otherwise $(x/y)^{2^{3k}+1}=(x/y)^{2^k+1}=1$ which yields that $(x/y)^{2^{2k}-1}=1$. Denote by $x=ty$. Since $\gcd(2k,n)=d'$, then $t\in\mathbb{F}_{2^{d'}}^*$.

- If $d'=d$, then $t\in\mathbb{F}_{2^d}^*$ and (14) is equivalent to $x^2+y^2=0$, that is, $x=y$ which has 2^n-1 solutions in $\mathbb{F}_q^*$. Therefore
$$M_2=1+(2^n-1)=2^n.$$
- If $d'=2d$, then by (9) we get (14) is equivalent to $x^{2^d+1}+y^{2^d+1}=0$. Then we have $t^{2^d+1}=1$ which has 2^d+1 solutions in $\mathbb{F}_{2^{2d}}^*$. Therefore
$$M_2=(2^d+1)(2^n-1)+1=2^{n+d}+2^n-2^d.$$

(iii). We have

$$\sum_{\alpha,\beta\in\mathbb{F}_q} T(\alpha,\beta)^3 = M_3\cdot 2^{2n} \quad \text{where}$$

$$M_3=\#\left\{(x,y,z)\in\mathbb{F}_q^3 \,\middle|\, x^{2^{3k}+1}+y^{2^{3k}+1}+z^{2^{3k}+1}=0, x^{2^k+1}+y^{2^k+1}+z^{2^k+1}=0\right\}$$
$$=M_2+T'\cdot(2^n-1) \tag{15}$$

and T' is the number of $\mathbb{F}_q$-solutions of

$$\begin{cases} x^{2^{3k}+1}+y^{2^{3k}+1}+1=0 \\ x^{2^k+1}+y^{2^k+1}+1=0. \end{cases} \tag{16}$$

Canceling y we have $\left(x^{2^{3k}+1}+1\right)^{2^k+1}=\left(x^{2^k+1}+1\right)^{2^{3k}+1}$ which is equivalent to

$$(x^{2^{4k}}+x)(x^{2^k}+x^{2^{3k}})=0.$$

Therefore $x^{2^{4k}}=x$ or $x^{2^k}=x^{2^{3k}}$.

- If $n/d \equiv 2 \pmod 4$, then $x \in \mathbb{F}_{2^{2d}}$ and symmetrically $y \in \mathbb{F}_{2^{2d}}$. Hence (16) is equivalent to $x^{2^d+1} + y^{2^d+1} + 1 = 0$ which is the well-known Hermitian curve on $\mathbb{F}_{2^{2d}}$. If follows that $T' = 2^{3d} - 2^d$.

- If $n/d \equiv 0 \pmod 4$, then $x \in \mathbb{F}_{2^{4d}}$ and hence $y \in \mathbb{F}_{2^{4d}}$. In this case $\left(x^{2^k+1} + y^{2^k+1} + 1\right)^{2^{3k}} = x^{2^{3k}+1} + y^{2^{3k}+1} + 1$ and then (16) is equivalent to $x^{2^d+1}+y^{2^d+1}+1 = 0$ which is a minimal curve on $\mathbb{F}_{2^{4d}}$ with genus $2^{d-1}(2^d-1)$. Hence $T' = 2^{4d} + 1 - 2^d(2^d-1)2^{2d} - (2^d+1) = 2^{3d} - 2^d$ (Note that there are exactly 2^d+1 rational infinite places on the curve $x^{2^d+1} + y^{2^d+1} + 1 = 0$ defined on $\mathbb{F}_q$).

Anyway, $M_3 = (2^{n+d} + 2^n - 2^d) + (2^n - 1)(2^{3d} - 2^d) = 2^{n+3d} + 2^n - 2^{3d}$.

(iv). We can calculate

$$\sum_{\alpha,\beta,\gamma \in \mathbb{F}_q} S(\alpha,\beta,\gamma)^3 = L_3 \cdot 2^{3n} \quad \text{where } L_3 \text{ is the number of solutions to}$$

$$x^{2^{3k}+1} + y^{2^{3k}+1} + z^{2^{3k}+1} = x^{2^k+1} + y^{2^k+1} + z^{2^k+1} = x + y + z = 0 \qquad (17)$$

- If $xyz = 0$, we may assume $x = 0$ and then $y = z$ which gives 2^n solutions. So is $y = 0$ or $z = 0$. Note that $x = y = z = 0$ has been counted 3 times. Hence there are exactly $3 \cdot 2^n - 2$ solutions to (17) satisfying $xyz = 0$.

- If $xyz = 0$, then L_3 is equal to $2^n - 1$ times the number of solutions to
$$x^{2^{3k}+1} + y^{2^{3k}+1} + 1 = x^{2^k+1} + y^{2^k+1} + 1 = x + y + 1 = 0 \qquad (18)$$
with $xy \neq 0$. Then we have $x^{2^{3k}+1} + (x+1)^{2^{3k}} + 1 = x^{2^k+1} + (x+1)^{2^k} + 1 = 0$ which is equivalent to $x^{2^{3k}} = x^{2^k} = x$. Hence $x \in \mathbb{F}_{2^{3k}} \cap \mathbb{F}_{2^k} \cap \mathbb{F}_{2^n}^* = \mathbb{F}_{2^d}^*$. Since $y \neq 0$, then $x \neq 1$. Therefore (18) has $2^d - 2$ solutions with $xy \neq 0$ and then $L_3 = 3 \cdot 2^n - 2 + (2^d - 2)(2^n - 1) = 2^{n+d} + 2^n - 2^d$.

Proof of Lemma 5: We get that

$$\begin{aligned}
qN &= \sum_{\omega \in \mathbb{F}_q} \sum_{x,y \in \mathbb{F}_q} (-1)^{\mathrm{Tr}_1^n\left(\omega\left(\alpha x^{2^{3k}+1} + \beta x^{2^k+1} + y^{2^d} + y\right)\right)} \\
&= q^2 + \sum_{\omega \in \mathbb{F}_q^*} \sum_{x \in \mathbb{F}_q} (-1)^{\mathrm{Tr}_1^n\left(\omega\left(\alpha x^{2^{3k}+1} + \beta x^{2^k+1}\right)\right)} \sum_{y \in \mathbb{F}_q} (-1)^{\mathrm{Tr}_1^n\left(y^{2^d}\left(\omega^{2^d} + \omega\right)\right)} \\
&= q^2 + q \sum_{\omega \in \mathbb{F}_{q_0}^*} \sum_{x \in \mathbb{F}_q} (-1)^{\mathrm{Tr}_1^n\left(\omega\left(\alpha x^{2^{3k}+1} + \beta x^{2^k+1}\right)\right)} \\
&= q^2 + q \sum_{\omega \in \mathbb{F}_{q_0}^*} T(\omega\alpha, \omega\beta)
\end{aligned}$$

where the 3-rd equality follows from that the inner sum is zero unless $\omega^{2^d} + \omega = 0$, i.e. $\omega \in \mathbb{F}_{q_0}$.

For any $\omega \in \mathbb{F}_{q_0}^*$, choose $t \in \mathbb{F}_{2^{2d}}$ such that $t^{2^d+1} = \omega$. Then $t^{2^{3k}+1} = t^{2^k+1} = \omega$. As a consequence $T(\omega\alpha, \omega\beta) = T(\alpha, \beta)$. Hence $N = 2^n + (2^d - 1) \cdot T(\alpha, \beta)$. $\square$

6 Appendix B

Proof of Theorem 1 (ii):

In the case n/d is even (that is, $d' = 2d$), we have $r_{\alpha,\beta} = s, s-2, s-4$ or $s-6$ for $(\alpha, \beta) \neq (0,0)$. According to Lemma 1 and Lemma 6, we get that if $r_{\alpha,\beta} = s - i$, then $T(\alpha, \beta) = (-1)^{m/d+\frac{i}{2}} 2^{\frac{n+id}{2}}$.

Combining Lemma 3 and Lemma 4 we have

$$n_0 + n_2 + n_4 + n_6 = 2^{2n} - 1 \tag{19}$$

$$n_0 - 2^d \cdot n_2 + 2^{2d} \cdot n_4 - 2^{3d} \cdot n_6 = (-1)^{m/d} 2^m (2^n - 1) \tag{20}$$

$$n_0 + 2^{2d} \cdot n_2 + 2^{4d} \cdot n_4 + 2^{6d} \cdot n_6 = 2^n (2^d + 1)(2^n - 1). \tag{21}$$

$$n_0 - 2^{3d} \cdot n_2 + 2^{6d} \cdot n_4 - 2^{9d} \cdot n_6 = (-1)^{m/d} 2^{m+3d} (2^n - 1). \tag{22}$$

Solving the system of equations consisting of (19)–(22) yields the result. $\square$

7 Appendix C

Proof of Theorem 2 : Define

$$\Xi = \left\{ (\alpha, \beta, \gamma) \in \mathbb{F}_q^3 \,|\, S(\alpha, \beta, \gamma) = 0 \right\}$$

and $\xi = \left|\Xi\right|$.

Recall $n_i, H_{\alpha,\beta}, , r_{\alpha,\beta}, A_\gamma$ in Section 1 and $N_{i,\varepsilon}, n_{i,\varepsilon},$ in the proof of Lemma 3.

- In the case n/d is odd, from Lemma 2, Lemma 3(i) and Lemma 4(iv) we have
$$n_1 + n_3 = 2^{2n} - 1 \tag{23}$$

$$n_1 + 2^{2d} \cdot n_3 = 2^{n-d}(2^d+1)(2^n-1). \tag{24}$$

These two equations yield

$$n_1 = \frac{(2^n-1)(2^{n+2d}-2^n-2^{n-d}+2^{2d})}{2^{2d}-1}, \quad n_3 = \frac{(2^{n-d}-1)(2^n-1)}{2^{2d}-1}.$$

From Lemma 2 we get that

$$\begin{aligned} \xi &= 2^n - 1 + (2^n - 2^{n-d})n_1 + (2^n - 2^{n-3d})n_3 \\ &= (2^n-1)(2^{2n} - 2^{2n-d} + 2^{2n-4d} + 2^n - 2^{n-d} - 2^{n-3d} + 1) \end{aligned} \tag{25}$$

- In the case n/d is even, from Lemma 2, Lemma 3(ii), Lemma 6 and Lemma 1(ii) we get that

$$\begin{aligned} \xi &= 2^n - 1 + (2^n - 2^{n-2d})n_{2,1} + (2^n - 2^{n-4d})n_{4,-1} + (2^n - 2^{n-6d})n_{6,1} \\ &= (2^n-1)\big[1 + (2^{2n} + 2^{2n-9d} - \varepsilon 2^{3m} + \varepsilon 2^{3m-d} + \varepsilon 2^{3m-3d} \\ &\quad -\varepsilon 2^{3m-5d} - \varepsilon 2^{3m-7d} + \varepsilon 2^{3m-8d} + 2^n - 2^{n-d} - 2^{n-4d} - 2^{n-6d})/(2^d+1)\big] \end{aligned} \tag{26}$$

By Lemma 2 we get the value distribution of $S(\alpha, \beta, \gamma)$ and then the weight distribution of $\mathcal{C}_2$. $\square$

Proof of Theorem 3 : For any possible value κ and $1 \le i, j \le 2$, define $M_\kappa(\mathcal{F}_i, \mathcal{F}_j)$ to be the frequency of κ in correlation values between any two sequences in $\mathcal{F}_i$ and $\mathcal{F}_j$ by any shift, respectively. Then the correlation distribution of sequences in $\mathcal{F}$ could be obtained if we can calculate all of the $M_\kappa(\mathcal{F}_i, \mathcal{F}_j)$. We will deal with it case by case.

- The correlation function between a_{α_1,β_1} and a_{α_2,β_2} by a shift τ $(0 \le \tau \le q-2)$ is

$$\begin{aligned} C_{(\alpha_1,\beta_1),(\alpha_2,\beta_2)}(\tau) &= \sum_{\lambda=0}^{q-2} (-1)^{a_{\alpha_1,\beta_1}(\lambda) - a_{\alpha_2,\beta_2}(\lambda+\tau)} \\ &= \sum_{\lambda=0}^{q-2} (-1)^{\mathrm{Tr}_1^n(\alpha_1 \pi^{\lambda(2^{3k}+1)} + \beta_1 \pi^{\lambda(2^k+1)} + \pi^\lambda) - \mathrm{Tr}_1^n(\alpha_2 \pi^{(\lambda+\tau)(2^{3k}+1)} + \beta \pi^{(\lambda+\tau)(2^k+1)} + \pi^{\lambda+\tau})} \\ &= S(\alpha', \beta', \gamma') - 1 \end{aligned}$$

where

$$\alpha' = \alpha_1 - \alpha_2 \pi^{\tau(2^{3k}+1)}, \quad \beta' = \beta_1 - \beta_2 \pi^{\tau(2^k+1)}, \quad \gamma' = 1 - \pi^\tau. \tag{27}$$

Fix $(\alpha_2,\beta_2)\in\mathbb{F}_q\times\mathbb{F}_q$, when (α_1,β_1) runs through $\mathbb{F}_q\times\mathbb{F}_q$ and τ takes values from 0 to $q-2$, (α',β',γ') runs through $\mathbb{F}_q\times\mathbb{F}_q\times\left\{\mathbb{F}_q\Big\backslash\{1\}\right\}$ exactly one time.

For any possible value κ of $S(\alpha,\beta,\gamma)$, define

$$s_\kappa=\#\left\{(\alpha,\beta,\gamma)\in\mathbb{F}_q\times\mathbb{F}_q\times\mathbb{F}_q\,|\,S(\alpha,\beta,\gamma)=\kappa+1\right\}\tag{28}$$

$$s_\kappa^1=\#\left\{(\alpha,\beta)\in\mathbb{F}_q\times\mathbb{F}_q\,\Big|\,S(\alpha,\beta,1)=\kappa+1\right\}\tag{29}$$

and

$$t_\kappa=\#\left\{(\alpha,\beta)\in\mathbb{F}_q\times\mathbb{F}_q\,|\,T(\alpha,\beta)=\kappa+1\right\}.\tag{30}$$

By Lemma 7 we have

$$s_\kappa^1=\frac{1}{2^n-1}\times(s_\kappa-t_\kappa).\tag{31}$$

Hence we get

$$M_\kappa(\mathcal{F}_1,\mathcal{F}_1)=2^{2n}\cdot\left(s_\kappa-s_\kappa^1\right)=2^{2n}\cdot\left(\frac{2^n-2}{2^n-1}\cdot s_\kappa+\frac{1}{2^n-1}\cdot t_\kappa\right).$$

- For the case n/d is odd. The cross correlation function between a_{α_1,β_1} and a_{α_2} by a shift τ $(0\le\tau\le q-2)$ is

  $$\begin{aligned}C_{(\alpha_1,\beta_1),\alpha_2}(\tau)&=\sum_{\lambda=0}^{q-2}(-1)^{a_{\alpha_1,\beta_1}(\lambda)-a_{\alpha_2}(\lambda+\tau)}\\&=\sum_{\lambda=0}^{q-2}(-1)^{\mathrm{Tr}_1^n(\alpha_1\pi^{\lambda(2^{3k}+1)}+\beta_1\pi^{\lambda(2^k+1)}+\pi^\lambda)-\mathrm{Tr}_1^n(\alpha_2\pi^{(\lambda+\tau)(2^{3k}+1)}+\pi^{(\lambda+\tau)(2^k+1)})}\\&=S(\alpha',\beta',1)-1\end{aligned}$$

  where $\alpha'=\alpha_1-\alpha_2\pi^{\tau(2^{3k}+1)},\beta'=\beta_1-\pi^{\tau(2^k+1)}$.

  Fix $0\le\tau\le q-2$ and $\alpha_2\in\mathbb{F}_q$, when (α_1,β_1) runs through $\mathbb{F}_q\times\mathbb{F}_q$, (α',β') runs through $\mathbb{F}_q\times\mathbb{F}_q$ exactly one time.

The cross correlation function between a_{α_1,β_1} and a by a shift τ $(0 \le \tau \le q-2)$ is

$$\begin{aligned} C_{(\alpha_1,\beta_1)}(\tau) &= \sum_{\lambda=0}^{q-2}(-1)^{a_{\alpha_1,\beta_1}(\lambda)-a(\lambda+\tau)} \\ &= \sum_{\lambda=0}^{q-2}(-1)^{\mathrm{Tr}_1^n(\alpha_1\pi^{\lambda(2^{3k}+1)}+\beta_1\pi^{\lambda(2^k+1)}+\pi^{\lambda})-\mathrm{Tr}_1^n(\pi^{(\lambda+\tau)(2^{3k}+1)})} \\ &= S(\alpha',\beta_1,1)-1 \end{aligned}$$

where $\alpha' = \alpha_1 - \pi^{\tau(2^{3k}+1)}$.

Fix $0 \le \tau \le q-2$, when (α_1,β_1) runs through $\mathbb{F}_q \times \mathbb{F}_q$, (α',β_1) runs through $\mathbb{F}_q \times \mathbb{F}_q$ exactly one time.

In total, by (28) and (29) we get

$$M_\kappa(\mathcal{F}_1,\mathcal{F}_2) = M_\kappa(\mathcal{F}_2,\mathcal{F}_1) = 2^n(2^n-1)\cdot s_\kappa^1 + (2^n-1)\cdot s_\kappa^1 = (2^n+1)(s_\kappa - t_\kappa).$$

- For the case n/d is odd. The cross correlation function between a_{α_1} and a_{α_2} by a shift τ $(0 \le \tau \le q-2)$ is

$$\begin{aligned} C_{\alpha_1,\alpha_2}(\tau) &= \sum_{\lambda=0}^{q-2}(-1)^{a_{\alpha_1}(\lambda)-a_{\alpha_2}(\lambda+\tau)} \\ &= \sum_{\lambda=0}^{q-2}(-1)^{\mathrm{Tr}_1^n(\alpha_1\pi^{\lambda(2^{3k}+1)}+\pi^{\lambda(2^k+1)})-\mathrm{Tr}_1^n(\alpha_2\pi^{(\lambda+\tau)(2^{3k}+1)}+\pi^{(\lambda+\tau)(2^k+1)})} \\ &= T(\alpha',\beta')-1 \end{aligned}$$

where $\alpha' = \alpha_1 - \alpha_2\pi^{\tau(2^{3k}+1)}, \beta' = 1-\pi^{\tau(2^k+1)}$.

When (α_1,α_2) runs through $\mathbb{F}_q \times \mathbb{F}_q$ and τ takes value from 0 to $q-2$, (α',β') runs through $\mathbb{F}_q \times \mathbb{F}_q\backslash\{1\}$ exactly q times.

The cross correlation function between a_{α_1} and a by a shift τ $(0 \le \tau \le q-2)$ is

$$\begin{aligned} C_{\alpha_1}(\tau) &= \sum_{\lambda=0}^{q-2}(-1)^{a_{\alpha_1}(\lambda)-a(\lambda+\tau)} \\ &= \sum_{\lambda=0}^{q-2}(-1)^{\mathrm{Tr}_1^n(\alpha_1\pi^{\lambda(2^{3k}+1)}+\pi^{\lambda(2^k+1)})-\mathrm{Tr}_1^n(\pi^{(\lambda+\tau)(2^{3k}+1)})} \\ &= T(\alpha',1)-1 \end{aligned}$$

where $\alpha' = \alpha_1 - \pi^{\tau(2^{3k}+1)}$. For fixed τ, $0 \le \tau \le q-2$, when α_1 runs through $\mathbb{F}_q$, then so is α'.

The auto-correlation function of a by a shift τ $(0 \le \tau \le q-2)$ is

$$\begin{aligned} C(\tau) &= \sum_{\lambda=0}^{q-2} (-1)^{a(\lambda)-a(\lambda+\tau)} \\ &= \sum_{\lambda=0}^{q-2} (-1)^{\mathrm{Tr}_1^n(\pi^{\lambda(2^{3k}+1)})-\mathrm{Tr}_1^n(\pi^{(\lambda+\tau)(2^{3k}+1)})} \\ &= T(1-\pi^{\tau(2^{3k}+1)}, 0) - 1. \end{aligned}$$

When τ takes values from 0 to $q-2$, then $1-\pi^{\tau(2^{3k}+1)}$ runs through $\mathbb{F}_q \backslash \{1\}$. Define

$$t_\kappa^0 = \#\{\beta \in \mathbb{F}_q |\, T(\alpha, 0) = \kappa+1\} \tag{32}$$

and

$$t_\kappa^1 = \#\{\beta \in \mathbb{F}_q |\, T(\alpha, 1) = \kappa+1\}. \tag{33}$$

Then a routine calculation shows that

$$t_\kappa^0 = \begin{cases} 1, & \kappa = 2^n - 1 \\ 2^n - 1, & \kappa = -1. \end{cases}$$

By Lemma 18 we have

$$t_\kappa^1 = \frac{1}{2^n-1}(t_\kappa - t_\kappa^0).$$

By Inclusion-Exclusion principle, Lemma 7, (30) and (32) we get

$$M_\kappa(\mathcal{F}_2, \mathcal{F}_2) = \left(\frac{2^n-2}{2^n-1}(t_\kappa - t_\kappa^0) + t_\kappa^0 \right) \cdot 2^n.$$

In total, sum up all the $M_\kappa(\mathcal{F}_i, \mathcal{F}_i)$ for $1 \le i, j \le 2$ and the result follows from Theorem 1 and Theorem 2.

□

8 Conclusion and Further Study

In this paper we have studied the exponential sums $T(\alpha, \beta)$ and $S(\alpha, \beta, \gamma)$ with $\alpha, \beta, \gamma \in \mathbb{F}_{2^n}$. After giving the value distribution of $T(\alpha, \beta)$ and $S(\alpha, \beta, \gamma)$, we determine the weight distributions of the cyclic codes $\mathcal{C}_1$ and $\mathcal{C}_2$ and the correlation distribution among a family of sequences.

In particular, for the case $f(x) = x^{2^{lk}+1} + x^{2^k+1}$ with $l \geq 4$, we could get the possible values of $\sum\limits_{x \in \mathbb{F}_q} (-1)^{\mathrm{Tr}_1^n(f(x))}$ and $\sum\limits_{x \in \mathbb{F}_q} (-1)^{\mathrm{Tr}_1^n(f(x)+\gamma x)}$. But the first three moment identities developed in Lemma 4 is not enough to determine the value distribution. However, we could get the possible weights of the corresponding cyclic codes. New machinery and technique should be invented to attack this problem.

In [13], the cross correlation distribution between two binary m-sequences $\{s_t\}$ and $\{s_{dt}\}$ with $d = \frac{2^{2k}+1}{2^k+1}$ for m odd and $k = 1$ has been determined. It seems that our techniques may be useful for the case $d = \frac{2^{lk}+1}{2^k+1}$ for a quite general l and k.

It turns out that the evaluation of $T(\alpha, \beta)$ and $S(\alpha, \beta, \gamma)$ are very similar for binary and odd characteristic when n/d is even. But the result is quite different when n/d is odd.

9 Acknowledgements

The authors will thank the anonymous referees for their helpful comments.

References

[1] A. Canteaut, P. Charpin, and H. Dobbertin, “Binary m-sequences with three-valued cross-correlation: a proof of Welchs conjecture,” *IEEE Trans. Inform. Theory,* vol. 46, no. 1, pp. 4–8, Jan. 2000.

[2] T. Cusick and H. Dobbertin, “Some new three-valued cross-correlation functions for binary m-sequences,” *IEEE Trans. Inform. Theory,* vol. 42, no. 4, pp. 1238–1240, April 1996.

[3] R.S. Coulter, “ON the evaluation of a class of Weil sums in characteristic 2,” *New Zeal. Jour. Math.,* vol. 28, pp. 171–184, 1999.

[4] C. Ding, T. Helleseth, and K.Y. Lam, “Several classes of binary sequences with three-level autocorrelation,” *IEEE Trans. Inf. Theory,* vol. 45, no. 7, pp. 2606–2612, Nov. 1999.

[5] C. Ding, T. Helleseth, and H. Martinsen, "New families of binary sequences with optimal three-valued autocorrelation," *IEEE Trans. Inf. Theory,* vol. 47, no. 1, pp. 428–433, Jan. 2001.

[6] H. Dobbertin, P. Felke, T. Helleseth, and P. Rosendahl, "Niho type cross-correlation functions via Dickson polynomials and Kloosterman sums," *IEEE Trans. Inform. Theory,* vol. 52, no. 2, pp. 613–627, Feb. 2006.

[7] R. Gold, "Maximal recursive sequences with 3-valued recursive cross-correlation functions," *IEEE Trans. Inf. Theory,* vol. 14, no. 1, pp. 154–156, Jan. 1968.

[8] T. Helleseth, "Some results about the cross-correlation function between two maximal linear sequences, " *Discrete Math.,* vol. 16, no. 3, pp. 209–232, 1976.

[9] T. Helleseth, "Pairs of m-sequences with a six-valued crosscorrelation," *Mathematical Mathematical Properties of Sequences and Other Combinatorial Structures,* J.-S. No, H.-Y. Song, T. Helleseth, and P. V. Kumar, Eds. pp. 1–6, Boston: Kluwer Academic Publishers, 2003.

[10] T. Helleseth, A. Kholosha, and G.J. Ness, "Characterization of m-Sequences of lengths $2^{2k} - 1$ and $2^k - 1$ with three-valued cross correlation," *IEEE Trans. Inform. Theory,* vol. 53, no. 6, pp. 2236–2245, June 2007.

[11] T. Helleseth and P.V. Kumar, "Sequences with low correlation," in *Handbook of Coding Theory*, V. S. Pless and W. C. Huffman, Eds. Amsterdam, The Netherlands: North-Holland, 1998.

[12] H. Hollmann and Q. Xiang, "A proof of Welch and Niho conjectures on cross-correlation of binary m-sequences," *Finite Fields and Their Appli.,* vol. 7, no. 2, pp. 253–286, 2001.

[13] A. Johansen and T. Helleseth, "A family of m-sequences with five-valued cross correlations", *IEEE Trans. Inf. Theory,* vol. 55, no. 2, pp. 880–887, Feb. 2009.

[14] J. Luo, S. Ling, C. Xing, "Cyclic codes and sequences from a class of Dembowski-Ostrom functions," preprint.

[15] J. Luo, Y. Tang, H. Wang "Exponential sums, cyclic codes and sequences: the binary Kasami case," preprint.

[16] T. Kasami, "Weight distribution of Bose-Chaudhuri-Hocquenghem codes," in Combinatorial Mathematics and Its Applications. R. C. Bose and T. A. Dowling, Eds. Chapel Hill, NC: Univerisity of North Carolina Press, 1969, pp. 335–357.

[17] T. Kasami, "Weight distribution formula for some class of cyclic codes," Coordinated Sci. Lab., Univ. Illinois, Urabana-Champaign, Tech. Rep. R-285(AD 635274), 1966.

[18] T. Kasami, "Weight enumerators for several classes of subcodes of the 2nd order Reed-Muller codes," *Inf. Contr.,* vol. 18, pp. 369–394, 1971.

[19] R. Lidl, and H. Niederreiter, Finite Fields, Addison-Wesley, Encyclopedia of Mathematics and its Applications, vol. 20, 1983.

[20] G. J. Ness and T. Helleseth, "A new family of four-valued cross correlation between m-sequences of different lengths," *IEEE Trans. Inf. Theory,* vol. 53, no. 11, pp. 4308–4313, Nov. 2007.

[21] G.J. Ness and T. Helleseth, "Cross correlation of m-sequences of different lengths," *IEEE Trans. Inf. Theory,* vol. 52, no. 4, pp. 1637–1648, Apri. 2006.

[22] Y. Niho, "Multivalued cross-correlation functions between two maximal linear recursive sequences," Ph.D dissertation, Univ. South.Calif., Los Angles, 1972.

[23] K.G. Paterson "Applications of exponential sums in communications theory," Extended Enterprise Laboratory, HP Laboratories Bristol, HPL-1999-101, 13th Sept., 1999.

[24] P. Rosendahl, "Niho type cross-correlation functions and related equations," Ph.D dissertation, Depart. Math., Univ. Turku, Finland, 2004.

[25] O.S. Rothaus, "Modified Gold codes," *IEEE Trans. Inf. Theory,* vol. 39, no. 2, pp. 654–656, Mar. 1993.

[26] M.K. Simon, J. Omura, R. Scholtz, and K. Levitt, "Spread Spectrum Communications," Rockville, MD: Computer Science, 1985, vol.ICIII.

[27] H.M. Trachtenberg, “On the cross-correlation function of maximal linear sequences,” Ph.D dissertation, Univ. South. Calif. Los Angles, 1970.

[28] M. Van Der Vlugt, “Surfaces and the weight distribution of a family of codes,” *IEEE Trans. Inf. Theory,* vol. 43, no. 4, pp. 1354–1360, Apri. 1997.

[29] J. Wolfmann, “Weight distribution of some binary primitive cylcic codes,” *IEEE Trans. Inf. Theory,* vol. 40, no. 6, pp. 2068–2071, Jun. 2004.

[30] Y. Xia, X. Zeng, and L. Hu, “The large set of p-ary Kasami sequences”, preprint.

[31] N.Y. Yu and G. Gong, “A new binary sequence family with low correlation and large size,”, *IEEE Trans. Inf. Theory,* vol. 52, no. 4, pp. 1624–1636, April 2006.

[32] N.Y. Yu and G. Gong, “New binary sequences with optimal autocorrelation magnitude,” *IEEE Trans. Inf. Theory,* vol. 54, no. 10, pp. 4771–4779, Oct. 2008.

[33] X. Zeng, L. Hu, W. Jiang, Q. Yue and X. Cao, “Weight distribution of a p-ary cyclic code,” see http://arxiv.org/abs/0901.2391, preprint.